

Answers to Even-Numbered Exercises

Exercises 1.1

2. (a) 2
(b) 1
(c) $f(2) = 1/2$
4. $x = 1$ and $x = 3$
10. (a) graph
(b) $\lim_{t \rightarrow 8^-} r(t) = 0.05$, $\lim_{t \rightarrow 8^+} r(t) = 0.10$
(c) all t except $t = 8$ and $t = 16$
12. 3
16. limit does not exist
20. 2
30. -6
32. $\frac{1}{6}$
40. 0
46. 1

Exercises 1.2

2. (a) 0 (b) $+\infty$ (c) $-\infty$ (d) 0
(e) $f(0) = -1$, $f(1) = 0$
8. $+\infty$
14. $+\infty$
20. $1/4$
32. 200
44. No vertical asymptote. Horizontal asymptote $y = 1$.
Always positive except at $x = 0$. Graph approaches $y = 1$ from below.

Exercises 1.3

4. -1, 4, 8

6. 1,6
8. 1
10. graph
22. (a) $f(1) = 1, f(2) = 4, f(3) = 6, f(4) = 2, f(5) = \sqrt{5}$,
(b) discontinuous at $x = 4$
28. continuous everywhere
30. $c = 12$
38. $-1/4$
42. f takes the value 2 twice and 3 once.
46. No. No; because the function is not continuous.
48. 4, they lies in: $(0.5,1)$ $(1,1.5)$ $(2.5,3)$ and $(3.5,4)$
50. (a) and (d)